My Pet's Health Passbook

Puppy & Kitten Vaccination and Deworming Record

Bring this to every vet visit. Ask your vet to write the date, stick the vaccine label and sign.

Dr. Vaishnavi Karmawat
B.V.Sc. & A.H.
Veterinarian & Author
vetvaishnavi.com

Pet's name: _____ Owner: _____

Dog / Cat • Breed: _____ My vet & phone: _____

Date of birth: _____ Sex: _____ Microchip / ID no.: _____

Puppy vaccination record

Age	Vaccine	Date given	Next due	Vet sign / label
6–8 weeks	1st puppy vaccine (DHPPi / Puppy DP) Distemper, Hepatitis, Parvo, Parainfluenza			
9–10 weeks	2nd vaccine (DHPPi + L, "7-in-1 / 9-in-1") Adds Leptospirosis			
12–13 weeks	3rd DHPPi + L • Anti-Rabies 1st dose			
15–16 weeks	Final puppy booster • Rabies booster if advised			
Every year	DHPPi + L booster • Anti-Rabies booster			
Optional	Kennel cough / Corona (ask your vet)			

Kitten vaccination record

Age	Vaccine	Date given	Next due	Vet sign / label
8–9 weeks	1st FVRCP (tricat) Cat flu (herpes, calici), Panleukopenia			
12 weeks	2nd FVRCP • Anti-Rabies 1st dose			
16 weeks	3rd FVRCP • Rabies booster if advised			
Every year	FVRCP booster • Anti-Rabies booster			

Good to know

- Keep puppies and kittens indoors, away from unknown animals, until 2 weeks after the last puppy/kitten shot.
- Mild sleepiness or soreness for a day after a vaccine is common.

Missed a dose? Adopted an adult?

- Don't skip it. Call your vet. A gap can mean the series needs restarting.
- Adult rescues with no record usually need 2 doses 3–4 weeks apart, plus rabies.

© Dr. Vaishnavi Karmawat • vetvaishnavi.com • For personal use only. Please don't share or resell.

Deworming & Tick-Flea Tracker

Dr. Vaishnavi Karmawat
B.V.Sc. & A.H.
Veterinarian & Author
vetvaishnavi.com

Worms and ticks are the most common reasons for weakness, poor growth and tick fever in Indian pets.
Your vet picks the dewormer and dose by your pet's current weight, so weigh before every dose.

Puppies

- **2 to 12 weeks:** every 2 weeks
- **3 to 6 months:** every month
- **Adults:** every 3 months

Kittens

- **3 to 8 weeks:** every 2 weeks
- **2 to 6 months:** every month
- **Adults:** every 3 months

Before each dose

- Weigh your pet.
- Use only the product your vet advised.
- Never use cattle or human dewormers.

Deworming log

Date	Age	Weight (kg)	Dewormer name	Dose given	Next due

Tick & flea prevention: 12-month tracker

write product & date

Jan	Feb	Mar	Apr	May	Jun

Jul	Aug	Sep	Oct	Nov	Dec

Tick check, every day in monsoon

- Run fingers over ears, between toes, neck and armpits.
- Remove ticks with a tick hook, never by squeezing.
- Clean bedding and corners of the house too.

See your vet fast if you notice

- Fever, not eating, very tired
- Pale gums, red spots on belly, nose bleed
- Dark or red urine

These can be signs of tick fever.

Disclaimer: Schedules shown are common guidelines in India. Your vet may change them based on your pet's health, the vaccine brand and local disease risk. Always follow your own veterinarian's advice.

© Dr. Vaishnavi Karmawat • vetvaishnavi.com • For personal use only. Please don't share or resell.